



Which vaccinations will my child receive?

Injection against RSV

- 0-2 weeks** for babies born from October through March
- 0-7 months** for babies born from April through September



6-9 weeks

- Rota
- (DTaP-IPV - Hib - HBV)*



3 months

- Rota
- DTaP-IPV - Hib - HBV
- Pneu



5 months

- DTaP-IPV - Hib - HBV
- Pneu



Abbreviations and what they mean

RS	Respiratory syncytial virus (RSV)
Rota	Rotavirus
D	Diphtheria
aP	Whooping cough (pertussis)
T	Tetanus
IPV	Polio
Hib	<i>Haemophilus influenzae</i> type b
HBV	Hepatitis B virus
Pneu	Pneumococcal disease
M	Mumps
M	Measles
R	Rubella
MenACWY	Meningococcal disease types ACWY
HPV	Human papillomavirus

Questions? Ask at the well-baby clinic or go to rijksvaccinatieprogramma.nl/en

12 months

- DTaP-IPV - Hib - HBV
- Pneu



14 months

- MMR
- MenACWY



3 years

- MMR



5 years

- DTaP



10 years

- HPV
- 2nd vaccination 6 months later



14 years

- DT-IPV
- MenACWY



* Only if the mother was not vaccinated against whooping cough during pregnancy (maternal whooping cough vaccination). This extra vaccine dose is also given in special circumstances. The paediatrician will discuss this with you.